

Contour Thread Lift Photo Gallery



WHO IS A CANDIDATE?

The Featherfacelift is an excellent office based procedure for patients with mild to moderate skin laxity and who want a noticeable improvement without any invasive surgery. Although the Featherfacelift results have been remarkable for many patients, it cannot replace the full effect of a face lift procedure.

POST-OPERATIVE

Some patients usually experience mild discomfort when smiling for approximately 1-2 weeks, although this usually resolves within a week. Post-operatively, patients will normally take Tylenol or other non-prescriptive analgesics for the mild discomfort that they usually will experience. Most patients can return to work within 48 hours. This is when the tape is usually removed. At this time, there will

be some light bruising usually present, and this can last anywhere from 2 days to 2 weeks. Typically, there is only a slight yellowish color to the skin and makeup may be applied immediately.

SCARS

There are tiny holes made in the skin from the insertion of the needle and typically these heal invisibly.

Post Procedure Photos





Fig 14: Preoperative and postoperative result 4 months after correction of the midface (4 sutures per side), neck (3 sutures per side) and brow (2 sutures per side)





Fig 15: Preoperative, 1 week postoperative and 6 month postoperative result. Patient with barbed suspension of midface & jowl (4 sutures per side), and neck (3 sutures per side). Note the initial “overcorrection” that should be noted in the first 1-2 weeks in order to achieve the desirable final result.





Fig 16: Preoperative and 2 month postoperative result after barbed suspension of the midface (4 sutures per side) and neck (3 sutures per side). Note prolonged postoperative facial edema that eventually resolved.





Fig 17 : Female patient preoperative and 4 month postoperative after barbed suspension of the midface area (4 sutures each side).



Fig 18: Preoperative and 7 month postoperative results after barbed suspension of the neck area only (4 sutures each side).